

Binding Energies and Scattering Observables in the $^4\text{He}_3$ Atomic System*

A. K. Motovilov^a, W. Sandhas^b, S. A. Sofianos^c, and E. A. Kolganova^{ad}

^aJINR, 141980 Dubna, Moscow Region, Russia

^bPI Universität Bonn, Endenicher Allee 11-13, D-53115 Bonn, Germany

^cPhysics Department, UNISA, P. O. Box 392, Pretoria 0003, South Africa

^dIAMS, Academia Sinica, PO Box 23-166, Taipei, Taiwan 10764, R. O. C.

The $^4\text{He}_3$ system is investigated using a hard-core version of the Faddeev differential equations and realistic ^4He – ^4He interactions. We calculate the binding energies of the ^4He trimer but concentrate in particular on scattering observables. The atom-diatom scattering lengths are calculated as well as the atom-diatom phase shifts for center of mass energies up to 2.45 mK.

There is a great number of experimental and theoretical studies of the ^4He three-atomic system (see, e. g., [1–9] and references cited therein). Most of the theoretical investigations consist merely in computing the bound states, while scattering processes found comparatively little attention. In Ref. [10] the characteristics of the He–He₂ scattering at zero energy were studied. The recombination rate of the reaction $(1 + 1 + 1 \rightarrow 2 + 1)$ was estimated in [11]. The phase shifts of the He–He₂ elastic scattering and breakup amplitudes at ultra-low energies have been calculated for the first time just recently [12] but only for the comparatively old HFD-B potential by Aziz *et al.* [13].

In the present paper we extend the investigations of Ref. [12]. There, the formalism, which consists of a hard-core version of the Faddeev differential equations, has been described in detail. As in [12] we use the finite-difference approximation of the two-dimensional partial-wave Faddeev equations. We consider only the case of zero total angular momentum and take $\hbar^2/m = 12.12 \text{ K } \text{\AA}^2$. In this work, we employ grids of the dimension 500–800 in both the hyperradius and hyperangle, while the cutoff hyperradius is chosen to be up to 1000 $\text{\AA}$. As compared to [12] we use in the present work the refined He–He interatomic potentials LM2M2 of Ref. [14], and TTY of Ref. [15]. Our numerical methods have also been substantially improved, which allowed us to deal with considerably larger grids. Furthermore, due to better computing facilities, we could take into account more partial waves.

Although we have performed detailed calculations of the $^4\text{He}_3$ binding energies, the main goal of this work was to perform calculations for the scattering of a helium atom off

*This work was supported by the Deutsche Forschungsgemeinschaft, Russian Foundation for Basic Research, and National Research Foundation of South Africa.
Contribution to Proceedings of 16th International Conference on Few-Body Problems in Physics, 6-10 March 2000, Taipei, Taiwan. LANL e-print physics/0009035.

Table 1

Dimer energies ϵ_d , ^4He – ^4He diatom scattering lengths $\ell_{\text{sc}}^{(2)}$, trimer ground-state energies $E_t^{(0)}$, trimer excited-state energies $E_t^{(1)}$, and ^4He atom – ^4He dimer scattering lengths ℓ_{sc} for the potentials used.

Potential	ϵ_d (mK)	$\ell_{\text{sc}}^{(2)}$ (Å)	l_{max}	$E_t^{(0)}$ (K)	$E_t^{(1)}$ (mK)	ℓ_{sc} (Å)
HFD-B	−1.68541	88.50	0	−0.0942	−2.45	168
			2	−0.1277	−2.71	138
			4	−0.1325	−2.74	135
LM2M2	−1.30348	100.23	0	−0.0891	−2.02	168
			2	−0.1213	−2.25	134
			4	−0.1259	−2.28	131
TTY	−1.30962	100.01	0	−0.0890	−2.02	168
			2	−0.1212	−2.25	134
			4	−0.1258	−2.28	131

a helium dimer at ultra-low energies. Our results for the trimer binding energies and ^4He –atom ^4He –dimer scattering lengths for the potentials employed are presented in Table 1. We also put in this table the corresponding dimer binding energies together with the ^4He – ^4He atomic scattering lengths. It should be noted that the main contribution to the trimer binding energies stems from the $l = 0$ and $l = 2$ partial-wave components, the latter being about 30 %, and is approximately the same for all potentials used. The contribution from the $l = 4$ partial wave is of the order of 3-4 % (cf. [5]). We notice that that our results for the ground-state energy $E_t^{(0)}$ of the trimer for $l_{\text{max}} = 4$ are in a perfect agreement with the corresponding values obtained in the most advanced calculations [4,7–9]. The same also holds true for the excited-state energy $E_t^{(1)}$. Our results for $E_t^{(1)}$ are in quite a good agreement with those of Refs. [8,9].

There are not many results in the literature concerning the He–He₂ scattering length. Apart from our previous result [12], there is that of Ref. [10] of $\ell_{\text{sc}} = 195$ Å, obtained within a zero-energy scattering calculation based on a separable approximation of the oldest Aziz *et al.* potential HFDHE2, and a more recent one obtained by Blume and Greene [16] via a Monte Carlo hyperspherical calculation with the LM2M2 potential. The latter result of $\ell_{\text{sc}} = 126$ Å is in good agreement with our result of 131 ± 5 Å (see Table 1). Within the accuracy of our calculations, the scattering lengths provided by the LM2M2 and TTY potentials, like the energies of the excited state, are exactly the same. It should be mentioned that in this case also the two-body binding energies and scattering lengths are almost identical.

We have also calculated the ^4He –atom ^4He –dimer scattering shifts for the HFD-B, LM2M2 and TTY potentials for center of mass energies up to 2.45 mK. After transformation to the laboratory system the phase shifts for the these potentials turn out to be practically the same, especially those for LM2M2 and TTY. Thus, in Fig. 1 we only plot the phase-shift results obtained for the HFD-B and TTY potentials. Note that for the phase shifts we use the normalization required by the Levinson theorem. Inclusion of the $l = 4$ partial-wave channel only adds about 0.5 % to the phase shifts obtained for $l = 0$ and $l = 2$. This is the reason why the corresponding curve for $l_{\text{max}} = 4$ is not depicted in Fig. 1.




Figure 1. ${}^4\text{He}$ atom – ${}^4\text{He}$ dimer scattering shifts for the HFD-B and TTY potentials. The lower curve corresponds to the case where $l_{\text{max}} = 0$ while for the upper $l_{\text{max}} = 2$.

A detailed exposition of the material presented, including tables for the phase shifts, is given in an extended paper [17].

REFERENCES

1. S. Grebenev, J. P. Toennies, and A. F. Vilesov, *Science* **279**, 2083 (1998).
2. F. Luo, C. F. Giese, and W. R. Gentry, *J. Chem. Phys.* **104**, 1151 (1996).
3. Th. Cornelius and W. Glöckle, *J. Chem. Phys.* **85**, 3906 (1986).
4. R. N. Barnett and K. B. Whaley, *Phys. Rev. A* **47**, 4082 (1993).
5. J. Carbonell, C. Gignoux, and S. P. Merkuriev, *Few-Body Systems* **15**, 15 (1993).
6. B. D. Esry, C. D. Lin, and C. H. Greene, *Phys. Rev. A* **54**, 394 (1996).
7. M. Lewerenz, *J. Chem. Phys.* **106**, 4596 (1997).
8. E. Nielsen, D. V. Fedorov, and A. S. Jensen, *J. Phys. B* **31**, 4085 (1998).
9. V. Roudnev and S. Yakovlev, LANL e-print physics/9910030.
10. S. Nakaichi-Maeda and T. K. Lim, *Phys. Rev. A* **28**, 692 (1983).
11. P. O. Fedichev, M. W. Reynolds, and G. V. Shlyapnikov, *Phys. Rev. Lett.* **77**, 2921 (1996).
12. E. A. Kolganova, A. K. Motovilov, and S. A. Sofianos, *J. Phys. B.* **31**, 1279 (1998).
13. R. A. Aziz, F. R. W. McCourt, and C. C. K. Wong, *Mol. Phys.* **61**, 1487 (1987).
14. R. A. Aziz and M. J. Slaman, *J. Chem. Phys.* **94**, 8047 (1991).
15. K. T. Tang, J. P. Toennies, and C. L. Yiu, *Phys. Rev. Lett.* **74**, 1546 (1995).
16. D. Blume and C. H. Green, *J. Chem. Phys.*, to appear.
17. A. K. Motovilov, W. Sandhas, S. A. Sofianos, and E. A. Kolganova, LANL e-print physics/9910016.